



January 18, 2013

Browse by: [Materials](#) | [Applications](#) | [Industries](#)[Terms](#) | [Submit News](#) | [Advertise](#) | [About](#)

Site Sponsors

Register today and find out the how and why of material characterization



Latest Nano News

[New Method to Amplify Signals in Graphene Oxide-Based Electrochemical Sensors](#)

[Mid-Infrared Frequency Combs Generated in Small Crystalline Micro-Resonators](#)

[UCLA to Develop New Nanoscale Materials for Semiconductor Components](#)

[New Pathway for Construction of Material Building Blocks at the Nanoscale](#)

[Atelier to Incorporate H₂O WaterBlock Nano-Coating in Luxury Smart Phones](#)

Site Sponsors

Posted in | [Nanoanalysis](#) | [Nanofabrication](#)

News Story

Recommend 0

Tweet 0

Share

0

Complex Nanomachine Mimics How Molecules Are Made in Nature

Published on January 11, 2013 at 5:30 AM

An industrial revolution on a minute scale is taking place in laboratories at The University of Manchester with the development of a highly complex machine that mimics how molecules are made in nature.

The artificial molecular machine developed by Professor David Leigh FRS and his team in the School of Chemistry is the most advanced molecular machine of its type in the world. Its development has been published in the journal *Science*.

Professor Leigh explains: "The development of this machine which uses molecules to make molecules in a synthetic process is similar to the robotic assembly line in car plants. Such machines could ultimately lead to the process of making molecules becoming much more efficient and cost effective. This will benefit all sorts of manufacturing areas as many manmade products begin at a molecular level. For example, we're currently modifying our machine to make drugs such as penicillin."

The machine is just a few nanometres long (a few millionths of a millimetre) and can only be seen using special instruments. Its creation was inspired by natural complex molecular factories where information from DNA is used to programme the linking of molecular building blocks in the correct order. The most extraordinary of these factories is the ribosome, a massive molecular machine found in all living cells.

Professor Leigh's machine is based on the ribosome. It features a functionalized nanometre-sized ring that moves along a molecular track, picking up building blocks located on the path and connecting them together in a specific order to synthesize the desired new molecule.

First the ring is threaded onto a molecular strand using copper ions to direct the assembly process. Then a "reactive arm" is attached to the rest of the machine and it starts to operate. The ring moves up and down the strand until its path is blocked by a bulky group. The reactive arm then detaches the obstruction from the track and passes it to another site on the machine, regenerating the active site on the arm. The ring is then free to move further along the strand until its path is obstructed by the next building block. This, in turn, is removed and passed to the elongation site on the ring, thus building up a new molecular structure on the ring. Once all the building blocks are removed from the track, the ring de-threads and the synthesis is over.

Professor Leigh says the current prototype is still far from being as efficient as the ribosome: "The ribosome can put together 20 building blocks a second until up to 150 are linked. So far we have only used our machine to link together 4 blocks and it takes 12 hours to connect each block. But you can massively parallel the assembly process: We are already using a million million million (10¹⁸) of these machines working in parallel in the laboratory to build molecules."

Professor Leigh continues: "The next step is to start using the machine to make sophisticated molecules with more building blocks. The potential is for it to be able to make molecules that have never been seen before. They're not made in nature and can't be made synthetically because of the processes currently used. This is a very exciting possibility for the future."

Source: <http://www.manchester.ac.uk/>Read in | [English](#) | [Español](#) | [Français](#) | [Deutsch](#) | [Português](#) | [Italiano](#) | [日本語](#) | [한국어](#) | [简体中文](#) | [繁體中文](#) | [Nederlands](#) | [Русский](#) | [Svenska](#)

Latest News

- [New Method to Amplify Signals in Graphene Oxide-Based Electrochemical Sensors](#)
- [Mid-Infrared Frequency Combs Generated in Small Crystalline Micro-Resonators](#)
- [UCLA to Develop New Nanoscale Materials for Semiconductor Components](#)
- [New Pathway for Construction of Material Building Blocks at the Nanoscale](#)